

Symmetry Properties of Scattering Amplitudes and Applications to Inverse Problems

A. G. RAMM

*Department of Mathematics, Kansas State University,
Manhattan, Kansas 66506*

Received September 14, 1989

For the transformations of the potential $q \rightarrow q \circ L$, where L is one of the following transformations: rotation, scaling, or translation, the corresponding transformation of the scattering amplitude is given. A necessary and sufficient condition on the scattering amplitude for the scatterer to be spherically symmetric is derived as an application of the above results. © 1991 Academic Press, Inc.

I. INTRODUCTION

Let

$$\begin{aligned} [\nabla^2 + k^2 - q(x)] u &= 0 \quad \text{in } \mathbb{R}^3, \quad k > 0, \\ u &= u_0(x, \theta, k) + A_q(\theta', \theta, k) r^{-1} \exp(ikr) + o(r^{-1}), \\ r = |x| &\rightarrow \infty, \quad x r^{-1} = \theta' \end{aligned} \quad (1)$$

$\theta, \theta' \in S^2$, the unit sphere, $u_0(x, \theta, k) := \exp(ik\theta \cdot x)$.

We are interested in the following question: if one changes $q(x)$ in (1) to $q \circ L$, what is the corresponding change in $A_q(\theta', \theta, k)$? For example, let

$$(q \circ L)(x) = (q \circ R)(x) := q(R^{-1}x), \quad R \in O(3), \quad (2)$$

where $O(3)$ is the group of rotations in \mathbb{R}^3 , or

$$(q \circ L)(x) = (q \circ \lambda)(x) := q(\lambda x), \quad \lambda > 0 \quad (3)$$

or

$$(q \circ L)(x) = (q \circ T_{-a})(x) := q(x - a), \quad a \in \mathbb{R}^3. \quad (4)$$

Here R , λ , and T_{-a} are operators of rotation, scaling by a factor $\lambda > 0$ and translation by a vector $-a$, respectively.

The basic results are the following formulas for the corresponding transformation of the scattering amplitudes:

$$A_q(\theta', \theta) = A_{q \circ R}(R\theta', R\theta), \quad (q \circ R)(x) := q(R^{-1}x) \quad (5)$$

$$A_q(\theta', \theta, k) = \lambda A_{\lambda^2 q \circ \lambda}(\theta', \theta, \lambda k), \quad (\lambda^2 q \circ \lambda)(x) = \lambda^2 q(\lambda x) \quad (6)$$

$$A_q(\theta', \theta, k) = A_{q \circ T_{-a}}(\theta', \theta, k) \exp\{ik(\theta' - \theta) \cdot a\}, \quad (q \circ T_{-a})(x) = q(x - a). \quad (7)$$

As an application, we prove that a necessary and sufficient condition for $q(x)$ to be spherically symmetric is that

$$A(\theta', \theta, k) = A(\theta' \cdot \theta, k), \quad \forall \theta', \theta \in S^2 \quad (8)$$

at a fixed $k > 0$ if

$$q \in Q_b := \{q : q = \bar{q}, q \in L^2(B_b), q = 0 \text{ for } |x| > b\}, \quad (9)$$

where $B_b := \{x : x \in \mathbb{R}^3, |x| \leq b\}$, $b > 0$ is an arbitrary fixed number and the overbar denotes complex conjugate. A similar result is proved for scattering by an obstacle. Namely, let

$$(\nabla^2 + k^2)u = 0 \quad \text{in } \Omega \subset \mathbb{R}^3, k > 0 \quad (10)$$

$$u_N + \zeta(s)u = 0 \quad \text{on } \Gamma, \operatorname{Im} \zeta(s) \geq 0 \quad (11)$$

$$u \text{ is of the form (2) with } A_{\Gamma, \zeta}(\theta', \theta, k) \text{ in place of } A_q(\theta', \theta, k). \quad (12)$$

Here Ω is the exterior of a bounded domain D with a smooth connected boundary Γ , N is the unit normal to Γ pointing into Ω .

Let us formulate the results (see also [10] and [11]).

THEOREM 1. *Formulas (5)–(7) hold.*

THEOREM 2. *Assume that (8) holds and $A(\theta', \theta, k) := A_q(\theta', \theta, k)$. Then (8) holds at a fixed $k > 0$ iff $q(x) = q(r)$.*

THEOREM 3. *Assume that (8) holds at a fixed $k > 0$ with $A(\theta', \theta, k) := A_{\Gamma, \zeta}(\theta', \theta, k)$. Then Γ is a sphere and $\zeta(s) = \text{const}$. The converse is trivially true.*

In Section II we give proofs. The results in Theorems 2 and 3 are based on formula (5) and the uniqueness theorems established in [1–4]. The ideas from [5, 6] are used. Our results cover the result in [7], where the two-dimensional scattering by obstacle with Dirichlet boundary condition has been studied, and the results in [8], where the scattering by an

obstacle with Dirichlet and Neumann boundary conditions has been treated and the scattering by a locally inhomogeneous acoustic medium has been treated. In the last case in [8] condition (8) is assumed to hold for all $k > 0$, while our Theorem 2 covers this case with the weaker assumption that (8) holds at a single $k > 0$. *It follows from our result that if (8) holds at a single $k > 0$ then (8) holds for all $k > 0$ (provided that (9) holds).*

II. PROOFS

1. Proof of Formulas (5)–(7)

Formulas (5)–(7) are direct consequences of the definition of the scattering amplitude.

Formula (5) means that the scattering amplitude is the same in a coordinate system τ and the coordinate system $R\tau$ in which each vector becomes Ra , where $R \in O(3)$ is an arbitrary rotation, and $O(3)$ is the group of rotations in \mathbb{R}^3 . The “rotated” potential $q \circ R$ is defined in (5).

A formal derivation of formulas (5)–(7) is based on writing the asymptotics of the scattering solution $u(x, \theta, k)$ in the new coordinate system. This derivation is the same in all three cases. Let us give it briefly.

(a) *Formula (5).* Consider

$$[\nabla_x^2 + k^2 - q(R^{-1}x)]u = 0 \quad \text{in } \mathbb{R}^3 \quad (13)$$

$$u(x, \theta, k) = \exp(ik\theta \cdot x) + A_{q \circ R}(\theta', \theta, k)|x|^{-1} \\ \times \exp(ik|x|) + o(|x|^{-1}), \quad \theta' = \frac{x}{|x|} := x^0. \quad (14)$$

Let $\xi := R^{-1}x$, $\xi^0 := \xi/|\xi| = R^{-1}\theta'$, $|\xi| = |x|$. Note that

$$R\theta' \cdot R\theta = \theta' \cdot \theta, \quad Rx \cdot a = x \cdot R^{-1}a \quad (14')$$

since $R' = R^{-1}$ for any $R \in O(3)$. Here R' is the transposed operator of rotation. Note that $\nabla_x^2 = \nabla_\xi^2$ is invariant under rotations. Write (13) and (14) in the ξ -coordinates:

$$[\nabla_\xi^2 + k^2 - q(\xi)]w = 0, \quad (15)$$

$$u = \exp(ik\theta \cdot R\xi) + A_{q \circ R}(\theta', \theta, k)|\xi|^{-1} \\ \times \exp(ik|\xi|) + o(|\xi|^{-1}), \quad \theta' = R\xi^0. \quad (16)$$

Note that, by (14), $\theta \cdot R\xi = R^{-1}\theta \cdot \xi$. Thus, the function (16) is the scattering solution

$$w = \exp(ikR^{-1}\theta \cdot \xi) + A_q(\alpha', R^{-1}\theta)|\xi|^{-1} \exp(ik|\xi|) + o(|\xi|^{-1}) \quad (16')$$

corresponding to the incident direction $R^{-1}\theta$. Let $R^{-1}\theta := \alpha$, $R^{-1}\theta' = \alpha' := \xi^0$. Then

$$A_q(\alpha', \alpha) = A_{q \circ R}(R\alpha', R\alpha), \quad \forall \alpha, \alpha' \in S^2.$$

This is formula (5).

(b) *Formula (6).* Consider

$$[\nabla_x^2 + k^2 - q(\lambda x)]u = 0 \quad \text{in } \mathbb{R}^3, \quad (17)$$

$$u = \exp(ik\theta \cdot x) + A_{q \circ \lambda}(\theta', \theta, k)|x|^{-1} \exp(ik|x|) + o(|x|^{-1}). \quad (18)$$

Let $\xi = \lambda x$. Then (17) can be written as

$$\left[\nabla_\xi^2 + \frac{k^2}{\lambda^2} - \frac{q(\xi)}{\lambda^2} \right] w = 0. \quad (19)$$

The scattering solution, corresponding to (19), is

$$w = \exp\left(i\frac{k}{\lambda}\theta \cdot \xi\right) + A_{\lambda^{-2}q}\left(\theta', \theta, \frac{k}{\lambda}\right) \frac{\exp(i(k/\lambda)|\xi|)}{|\xi|} + o\left(\frac{1}{|\xi|}\right) \quad (20)$$

while (18) can be written as

$$u = \exp\left(i\frac{k}{\lambda}\theta \cdot \xi\right) + \lambda A_{q \circ \lambda}(\theta', \theta, k)|\xi|^{-1} \exp\left(i\frac{k}{\lambda}|\xi|\right) + o\left(\frac{1}{|\xi|}\right). \quad (21)$$

Compare (19), (20), and (21) to obtain

$$A_{\lambda^{-2}q}\left(\theta', \theta, \frac{k}{\lambda}\right) = \lambda A_{q \circ \lambda}(\theta', \theta, k). \quad (22)$$

Let $k/\lambda := \mu$, $\lambda^{-2}q = p$. Then (22) can be written as

$$A_p(\theta', \theta, \mu) = \lambda A_{\lambda^2 p \circ \lambda}(\theta', \theta, \lambda\mu). \quad (23)$$

This is formula (6) with $p = q$ and $\mu = k$.

(c) *Formula (7).* Consider

$$[\nabla_x^2 + k^2 - q(x - a)]u = 0 \quad \text{in } \mathbb{R}^3 \quad (24)$$

$$u = \exp(ik\theta \cdot x) + A_{q \circ T_{-a}}(\theta', \theta, k)|x|^{-1} \exp(ik|x|) + o(|x|^{-1}). \quad (25)$$

Let $x = a = \xi$. Then (24) becomes

$$[\nabla_x^2 + k^2 - q(\xi)]w = 0, \quad (26)$$

and

$$w = \exp(ik\theta \cdot \xi) + A_q(\theta', \theta, k) \frac{\exp(ik|\xi|)}{|\xi|} + o\left(\frac{1}{|\xi|}\right) \quad (27)$$

while

$$\begin{aligned} u &= \exp(ik\theta \cdot \xi) \exp(ik\theta \cdot a) + A_{q \cdot T_{-a}}(\theta', \theta, k) |\xi + a|^{-1} \\ &\quad \times \exp(ik|\xi + a|) + o\left(\frac{1}{|\xi|}\right). \end{aligned} \quad (28)$$

Note that

$$\begin{aligned} |\xi + a| &= |\xi| + \xi^0 \cdot a + O(|\xi|^{-1}) \quad \text{as } |\xi| \rightarrow \infty, \\ \xi^0 &:= \frac{\xi}{|\xi|} \rightarrow x^0 = \theta' \quad \text{as } |\xi| \rightarrow \infty. \end{aligned} \quad (29)$$

Thus (28) becomes

$$\begin{aligned} u &= \exp(ik\theta \cdot a) \left[\exp(ik\theta \cdot \xi) + A_{q \cdot T_{-a}}(\theta', \theta, k) \right. \\ &\quad \left. \times \exp\{ik(\theta' - \theta) \cdot a\} \frac{\exp(ik|\xi|)}{|\xi|} + o\left(\frac{1}{|\xi|}\right) \right]. \end{aligned} \quad (30)$$

Since u solves the homogeneous linear equation (24) the expression in brackets in (30) solves Eq. (26). Compare (27) and (30) to obtain

$$A_q(\theta', \theta, k) = A_{q \cdot T_{-a}}(\theta', \theta, k) \exp\{ik(\theta' - \theta) \cdot a\}. \quad (31)$$

This is formula (7).

2. Proof of Theorems 2 and 3

(a) *Proof of Theorem 2.* The basic auxiliary result is the following uniqueness theorem from [3, 4].

PROPOSITION 1. *Let $q_j \in Q_b$, $j = 1, 2$. If $A_{q_1}(\theta', \theta, k) = A_{q_2}(\theta', \theta, k)$ at a fixed $k > 0$ and all $\theta', \theta \in S^2$, then $q_1 = q_2$.*

Theorem 2 follows immediately from formula (5) and Proposition 1.

Indeed, it is well known (and follows immediately from separation of variables) that if the potential is spherically symmetric

$$q(x) = q(|x|) = q(r), \quad r = |x| \quad (32)$$

then (8) holds for all $k > 0$. Assume that (9) holds and (8) holds at a fixed $k > 0$ with $A = A_q$. Then

$$\begin{aligned} A_q(\theta', \theta, k) &= A_q(\theta' \cdot \theta, k) = A_q(R\theta' \cdot R\theta, k) \\ &= A_{q \circ R}(R\theta', R\theta, k), \quad \forall R \in O(3). \end{aligned} \quad (33)$$

Here the second equation follows from (14) and the third equation is formula (5). Since $q \circ R \in Q_b$ if $q \in Q_b$, and since θ' and θ are arbitrary, one can write (33) as

$$A_q(\alpha', \alpha, k) = A_{q \circ R}(\alpha', \alpha, k), \quad \forall R \in O(3), \forall \alpha', \alpha \in S^2, \quad (34)$$

where $\alpha' = R\theta'$, $\alpha = R\theta$. By Proposition 1, it follows that

$$q = q \circ R, \quad \forall R \in O(3). \quad (35)$$

This is equivalent to (32). Theorem 2 is proved.

(b) *Proof of Theorem 3.* The basic auxiliary result is the following uniqueness theorem from [1, p. 85].

PROPOSITION 2. Assume that $A_{\Gamma_1, \zeta_1}(\theta', \theta, k) = A_{\Gamma_2, \zeta_2}(\theta', \theta, k)$ for all $\theta', \theta \in S^2$ and a fixed $k > 0$. Then $\Gamma_1 = \Gamma_2$ and $\zeta_1(s) = \zeta_2(s)$.

Note that if Γ is a sphere and $\zeta = \text{const}$, then (8) holds with $A(\theta', \theta, k) = A_{\Gamma, \zeta}(\theta', \theta, k)$ for all $\theta', \theta \in S^2$ and all $k > 0$. This follows from the analytical solution of the scattering problem by separation of variables. Assume now that (8) holds at a single fixed $k > 0$. Then, by (8) and (14),

$$A_{\Gamma, \zeta}(\theta', \theta, k) = A_{\Gamma, \zeta}(\theta' \cdot \theta, k) = A_{\Gamma, \zeta}(R\theta' \cdot R\theta, k) \quad (36)$$

and

$$A_{\Gamma, \zeta}(\theta', \theta, k) = A_{\Gamma \circ R, \zeta \circ R}(R\theta' \cdot R\theta, k) \quad (37)$$

by formula analogous to (5). Here $\Gamma \circ R$ is the surface Γ rotated by the element $R \in O(3)$, and $(\zeta \circ R)(s) = \zeta(R^{-1}s)$. From (36) and (37) it follows that

$$A_{\Gamma, \zeta}(\alpha', \alpha, k) = A_{\Gamma \circ R, \zeta \circ R}(\alpha', \alpha, k), \quad \forall R \in O(3), \forall \alpha', \alpha \in S^2. \quad (38)$$

By Proposition 2, Eq. (38) implies

$$\Gamma = \Gamma \circ R, \quad \zeta = \zeta \circ R, \quad \forall R \in O(3). \quad (39)$$

Thus Γ is a sphere and $\zeta(s) = \zeta(|s|) = \text{const}$ on the sphere Γ . Theorem 3 is proved.

Remark 1. Consider compactly supported potentials $q = q(\rho, z)$, $z = x_3$, $\rho = (x_1^2 + x_2^2)^{1/2}$. Let R_φ , $0 \leq \varphi \leq 2\pi$, denote rotations about the x_3 -axis. Then $q \circ R_\varphi = q$. By formula (5) for $R = R_\varphi$ one has

$$A_q(\theta', \theta, k) = A_{q \circ R_\varphi}(R_\varphi \theta', R_\varphi \theta, k) = A_q(R_\varphi \theta', R_\varphi \theta, k). \quad (40)$$

This symmetry property

$$A_q(\theta', \theta, k) = A_q(R_\varphi \theta', R_\varphi \theta, k) \quad (41)$$

is a necessary property of the scattering amplitude corresponding to the potential which is axially symmetric about the x_3 -axis; i.e., $q(x) = q(\rho, z)$. If $q \in Q_b$, then (41) is also sufficient for $q(x)$ to be axially symmetric about the x_3 -axis. This is proved as in Theorem 1.

Remark 2. Consider the equation

$$[\nabla^2 + k^2 + k^2 v(x)] u = 0 \quad \text{in } \mathbb{R}^3, \quad k > 0, \quad (42)$$

for $v \in Q_b$. At a fixed $k > 0$ the scattering of a plane wave by the inhomogeneity $v(x)$ is identical with the scattering of this wave by the potential $q(x) = -k^2 v(x)$. Therefore, by Theorem 2, $v(x) = v(|x|)$ if and only if condition (8) holds at a fixed $k > 0$, where $A(\theta', \theta, k)$ is the scattering amplitude corresponding to Eq. (42).

Remark 3. The argument in this paper is based on uniqueness theorems. If $q \in Q^{(\beta)} := \{q: |q(x)| \leq c(1 + |x|)^{-\beta}, \quad q = \bar{q}\}$, $\beta > 3$, Proposition 2 is no longer valid. However, for this class of potentials the following *uniqueness theorem* holds: if $A_{q_1}(\theta', \theta, k) = A_{q_2}(\theta', \theta, k)$ for all $\theta', \theta \in S^2$ and almost all $k > 0$, then $q_1 = q_2$. The argument used in the proof of Theorem 2 shows that if $q \in Q^{(\beta)}$, $\beta > 3$, and if (8) holds for all $k > 0$, then $q(x) = q(|x|)$. The uniqueness theorem holds even for $q \in Q^{(\beta)}$ with $\beta > 1$ (see [9]). Therefore, the following theorem holds.

THEOREM 4. If $q \in Q^{(\beta)}$, $\beta > 1$, then $q(x) = q(r)$ iff (8) holds for almost all $k > 0$.

Remark 4. If $q \in Q^{(\beta)}$, $1 < \beta \leq 3$, then $A(\theta', \theta, k)$ is not necessarily continuous in θ' and θ so that (8) is understood as equality of kernels of operators on $L^2(S^2)$.

ACKNOWLEDGMENT

The author thanks NSF, ONR, and SFB 256 for support and Professor R. Leis for a discussion. This paper was written while the author was a guest of SFB 256 at the University of Bonn.

REFERENCES

1. A. G. RAMM, "Scattering by Obstacles," Reidel, Dordrecht, 1986.
2. A. G. RAMM, Multidimensional inverse problems and completeness of the products of solutions to Partial Differential Equations, *J. Math. Anal. Appl.* **134** (1988), 211–253.
3. A. G. RAMM, Uniqueness theorems for multidimensional inverse problems with unbounded coefficients, *J. Math. Anal. Appl.* **136** (1988), 568–574.
4. A. G. RAMM, Recovery of the potential from fixed energy scattering data, *Inverse Problems* **4** (1988), 877–886.
5. A. G. RAMM, Necessary and sufficient condition for a scattering amplitude to correspond to a spherically symmetric scatterer, *Appl. Math. Lett.* **2** (1989), 263–265.
6. A. G. RAMM AND O. L. WEAVER, Necessary and sufficient condition on the fixed energy scattering data for the potential to be spherically symmetric, *Inverse Problems* **5** (1989), L45–47.
7. S. KARP, Far field amplitudes and inverse diffraction theory, in "Electromagnetic Waves" (R. Langer, Ed.), pp. 291–300, Univ. of Wisconsin Press, Madison, 1962.
8. D. COLTON AND A. KIRSCH, Karp's theorem in acoustic scattering theory, *Proc. Amer. Math. Soc.* **103** (1988), 783–788.
9. Y. SAITO, An asymptotic behavior of the S -matrix and the inverse scattering problem, *J. Math. Phys.* **21** (1984), 3105–3111.
10. A. G. RAMM, Completeness of the set of the products of solutions to PDE and inverse problems, *Inverse Problems* **6** (1990), 643–664.
11. A. G. RAMM, "Multidimensional Inverse Scattering Problems," Longman/Wiley, New York, 1992.